



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 1/21, 5/10, 15/12, 15/63</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/40698</b> <b>(43) International Publication Date:</b> 13 July 2000 (13.07.00)
<b>(21) International Application Number:</b> PCT/US98/27862 <b>(22) International Filing Date:</b> 31 December 1998 (31.12.98) <b>(71) Applicant (for all designated States except US):</b> THE GENERAL HOSPITAL CORPORATION [US/US]; Fruit Street, Boston, MA 92114 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> GARDELLA, Thomas, J. [US/US]; 1045 Greendale Avenue, Needham, MA 02492 (US). KRONENBERG, Henry, M. [US/US]; 48 Hastings Road, Belmont, MA 02178 (US). POTTS, John, T., Jr. [US/US]; 129 Chestnut Street, Newton, MA 02165 (US). <b>(74) Agents:</b> GOLDSTEIN, Jorge, A. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> <i>With an indication in relation to deposited biological material furnished under Rule 13bis separately from the description.</i> <i>With a statement concerning non-prejudicial disclosure or exception to lack of novelty.</i>
<b>(54) Title:</b> PTH RECEPTOR AND SCREENING ASSAY UTILIZING THE SAME		
<b>(57) Abstract</b>  The invention provides a novel PTH receptor polypeptide, r6Nt, characterized by a deletion of the extracellular amino-terminus, ligand binding domain of the receptor. Additionally disclosed are nucleic acid molecules encoding the receptor. The receptor has a minimal domain for ligand binding and is useful in screening assays designed for the identification of agonists and antagonists of PTH receptor activity.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

<b>AL</b>	Albania	<b>ES</b>	Spain	<b>LS</b>	Lesotho	<b>SI</b>	Slovenia
<b>AM</b>	Armenia	<b>FI</b>	Finland	<b>LT</b>	Lithuania	<b>SK</b>	Slovakia
<b>AT</b>	Austria	<b>FR</b>	France	<b>LU</b>	Luxembourg	<b>SN</b>	Senegal
<b>AU</b>	Australia	<b>GA</b>	Gabon	<b>LV</b>	Latvia	<b>SZ</b>	Swaziland
<b>AZ</b>	Azerbaijan	<b>GB</b>	United Kingdom	<b>MC</b>	Monaco	<b>TD</b>	Chad
<b>BA</b>	Bosnia and Herzegovina	<b>GE</b>	Georgia	<b>MD</b>	Republic of Moldova	<b>TG</b>	Togo
<b>BB</b>	Barbados	<b>GH</b>	Ghana	<b>MG</b>	Madagascar	<b>TJ</b>	Tajikistan
<b>BE</b>	Belgium	<b>GN</b>	Guinea	<b>MK</b>	The former Yugoslav Republic of Macedonia	<b>TM</b>	Turkmenistan
<b>BF</b>	Burkina Faso	<b>GR</b>	Greece	<b>ML</b>	Mali	<b>TR</b>	Turkey
<b>BG</b>	Bulgaria	<b>HU</b>	Hungary	<b>MN</b>	Mongolia	<b>TT</b>	Trinidad and Tobago
<b>BJ</b>	Benin	<b>IE</b>	Ireland	<b>MR</b>	Mauritania	<b>UA</b>	Ukraine
<b>BR</b>	Brazil	<b>IL</b>	Israel	<b>MW</b>	Malawi	<b>UG</b>	Uganda
<b>BY</b>	Belarus	<b>IS</b>	Iceland	<b>MX</b>	Mexico	<b>US</b>	United States of America
<b>CA</b>	Canada	<b>IT</b>	Italy	<b>NE</b>	Niger	<b>UZ</b>	Uzbekistan
<b>CF</b>	Central African Republic	<b>JP</b>	Japan	<b>NL</b>	Netherlands	<b>VN</b>	Viet Nam
<b>CG</b>	Congo	<b>KE</b>	Kenya	<b>NO</b>	Norway	<b>YU</b>	Yugoslavia
<b>CH</b>	Switzerland	<b>KG</b>	Kyrgyzstan	<b>NZ</b>	New Zealand	<b>ZW</b>	Zimbabwe
<b>CI</b>	Côte d'Ivoire	<b>KP</b>	Democratic People's Republic of Korea	<b>PL</b>	Poland		
<b>CM</b>	Cameroon	<b>KR</b>	Republic of Korea	<b>PT</b>	Portugal		
<b>CN</b>	China	<b>KZ</b>	Kazakstan	<b>RO</b>	Romania		
<b>CU</b>	Cuba	<b>LC</b>	Saint Lucia	<b>RU</b>	Russian Federation		
<b>CZ</b>	Czech Republic	<b>LI</b>	Liechtenstein	<b>SD</b>	Sudan		
<b>DE</b>	Germany	<b>LK</b>	Sri Lanka	<b>SE</b>	Sweden		
<b>DK</b>	Denmark	<b>LR</b>	Liberia	<b>SG</b>	Singapore		
<b>EE</b>	Estonia						

## PTH Receptor and Screening Assay Utilizing the Same

### *Background of the Invention*

#### *Statement as to Rights to Inventions Made Under Federally-Sponsored Research and Development*

5           Part of the work performed during development of this invention utilized U.S. Government funds. The U.S. Government has certain rights in this invention.

#### *Field of the Invention*

          The present invention is related to the fields of molecular biology, developmental biology, physiology, neurobiology, endocrinology and medicine.

#### *Related Art*

10           PTH is the principal regulator of blood calcium levels and mediates this action through binding to PTH-1 receptors on bone and kidney cells (Kronenberg, H.M., *et al.*, in "Handbook of Experimental Pharmacology, Springer-Verlag," Heidelberg (1993)). This receptor also responds to PTH-related peptide, a factor  
15           which plays a role in embryonic bone development and is the causative agent of hypercalcemia of malignancy (Lanske, B., *et al.*, *Science* 273:663-666 (1996)). PTH and PTHrP peptides have been shown to have potent anabolic effects on bone, and it is possible, therefore, that PTH-1 receptor agonists could ultimately be used to treat metabolic bone diseases, such as osteoporosis (Dempster, D.W.,  
20           *et al.*, *Endocr Rev.* 14(6):690-709 (1994)).

          In the fully bioactive PTH(1-34) peptide, the major determinants of receptor-binding affinity reside within amino acids 15 to 34 (Nussbaum, S.R., *et al.*, *J. Biol. Chem.* 255:10183-10187 (1980); Gardella, T.J., *et al.*, *Endocrinology*

132(5):2024-2030 (1993); Caulfield, M.P., *et al.*, *Endocrinology* 127:83-87 (1990); Abou-Samra, A. B., *et al.*, *Endocrinology* 125:2215-2217 (1989)), which are moderately conserved among PTHs and PTHrPs from various species (Suva, L.J., *et al.*, *Science* 237(4817):893-896 (1987)). The determinants of receptor activation lie within the more stringently conserved amino-terminal residues, and deletion of these residues yields competitive PTH-1 receptor antagonists (Horiuchi, N., *et al.*, *Science* 220:1053-1055 (1983); Nutt, R.F., *et al.*, *Endocrinology* 127:491-493 (1990)). Amino-terminal PTH or PTHrP fragments shorter in length than PTH(1-27) have not previously been found to be biologically active (Rosenblatt, M., *Pathobiology Annual*, Raven Press, New York, 11:53-84 (1981); Azarani, A., *et al.*, *J. Biol. Chem.* 271(25):14931-14936 (1996); Tregear, G.W., *et al.*, *Endocrinology* 93:1349-1353 (1973)), yet the functional importance and evolutionary conservation of the amino-terminal residues predicts that they directly interact with the receptor.

The PTH-1 receptor couples strongly to the adenylyl cyclase/protein kinase A signaling pathway and, in some settings, to other pathways including those mediated by phospholipase C/protein kinase C and intracellular calcium (About-Samra, A. B., *et al.*, *Endocrinology* 129:2547-2554 (1991); Jüppner, H., *et al.*, *Science* 254:1024-1026 (1991); Guo, J. *et al.*, *Endocrinology* 136:3884-3891 (1995); Hruska, K.A., *et al.*, *J. Clin. Invest.* 79:230-239 (1987); Donahue, H.J., *et al.*, *J. Biol. Chem.* 263:13522-13527 (1988)). The PTH-1 receptor is a member of the family B subgroup of G protein-coupled receptors, which also includes the receptors for calcitonin and secretin (Kolakowski, L. F., "GCRDb: A G-Protein-Coupled Receptor Database," *Receptors and Channels* 2:1-7 (1994)). Mutagenesis and crosslinking studies have indicated that multiple domains of these receptors contribute to ligand interaction, including the large amino-terminal extracellular domain, the extracellular loops and the transmembrane helices (Jüppner, H., *et al.*, *Endocrinology* 134:879-884 (1994); Lee, C., *et al.*, *Mol. Endo.* 9:1269-1278 (1995); Turner, P., *et al.*, *J. Bone Min. Res.* 12(1):Abstract 121 (1997); Dautzenberg, F., *et al.*, *Proc. Natl. Acad. Sci.*

95:4941-4946 (1998); Holtmann, M., *et al.*, *J. Biol. Chem.* 270:14394-14398 (1995); DeAlmeida, V. and Mayo, K., *Mol. Endo.* 12:750-765 (1998); Stroop, S., *et al.*, *Biochem.* 34:1050-1057 (1994); Zhou, A., *et al.*, *Proc. Natl. Acad. Sci. USA* 94:3644-3649 (1997); Bisello, A., *et al.*, *J. Biol. Chem.* 273:22498-22505 (1998)). Studies using PTH/calcitonin chimeric receptors and hybrid ligands have suggested a general topology of the interaction in which the amino-terminal extracellular domain of the receptor recognizes the carboxyl-terminal binding domain of the ligand, while the "core" region of the receptor containing the seven transmembrane helices and connecting loops recognizes the amino-terminal signaling portion of the ligand (Bergwitz, C., *et al.*, *J. Biol. Chem.* 271:26469-26472 (1996)). Similar conclusions were derived from earlier receptor chimera studies (Jüppner, H., *et al.*, *Endocrinology* 134:879-884 (1994); Stroop, S., *et al.*, *Biochem.* 34:1050-1057 (1994); Gardella, T.J., *et al.*, *Endocrinology* 135:1186-1194 (1994)) and from recent crosslinking studies with photoreactive PTH analogs (Bisello, A., *et al.*, *J. Biol. Chem.* 273:22498-22505 (1998); Mannstadt, M., *et al.*, *J. Biol. Chem.* 273:16890-16896 (1998)).

In the current study we investigate the signaling component of the interaction between PTH and the PTH-1 receptor using a domain-based approach. This approach employs short amino-terminal PTH fragment analogs and a PTH receptor mutant that lacks most of the amino-terminal extracellular domain. The results of cAMP-signaling assays performed with these smaller ligands and receptors demonstrate that the conserved amino-terminal (Kronenberg, H.M., *et al.*, in "Handbook of Experimental Pharmacology, Springer-Verlag," Heidelberg (1993); Lanske, B., *et al.*, *Science* 273:663-666 (1996); Dempster, D.W., *et al.*, *Endocr Rev.* 14(6):690-709 (1994); Nussbaum, S.R., *et al.*, *J. Biol. Chem.* 255:10183-10187 (1980); Gardella, T.J., *et al.*, *Endocrinology* 132(5):2024-2030 (1993); M. P. Caulfield *et al.*, *Endocrinology* 127:83-87 (1990); A. B. Abou-Samra *et al.*, *Endocrinology* 125:2215-2217 (1989); Suva, L.J., *et al.*, *Science* 237(4817):893-896 (1987); Horiuchi, N., *et al.*, *Science* 220:1053-1055 (1983); Nutt, R. F., *et al.*, *Endocrinology* 127:491-493 (1990); Rosenblatt, M.,

5 *Pathobiology Annual*, Raven Press, New York, 11:53-84 (1981); Azarani, A., *et al.*, *J. Biol. Chem.* 271(25):14931-14936 (1996); Tregear, G.W., *et al.*, *Endocrinology* 93:1349-1353 (1973); About-Samra, A.B., *et al.*, *Endocrinology* 129:2547-2554 (1991)) segment of PTH functions as an autonomous signaling domain and that this domain interacts with the core region of the receptor.

### ***Summary of the Invention***

10 PTH is the principal regulator of blood calcium levels and mediates this action through binding to PTH-1 receptors on bone and kidney cells. PTH-1 receptor agonists may ultimately be used to treat metabolic bone diseases, such as osteoporosis. Thus there is a strong need in the art develop new and improved PTH and PTH receptor reagents for the treatment of human disease.

15 In a first embodiment, the invention provides a novel PTH-1 receptor polypeptide, r $\delta$ Nt, characterized by a deletion of the extracellular amino-terminal ligand binding domain. The invention also provides nucleic acid molecules encoding the r $\delta$ Nt receptor polypeptide.

20 In a second embodiment the r $\delta$ Nt receptor of the invention is useful for screening procedures designed to identify agonists and antagonists of PTH receptor function. The invention provides screening utilizing either cAMP accumulation or competitive binding for the evaluation of test compounds with cells expressing the r $\delta$ Nt receptor.

### ***Brief Description of the Figures***

Figure 1. Presentation of the nucleic acid sequence and the amino acid sequence of the mutant PTH1R receptor, r $\delta$ Nt.

25 Figure 2. cAMP-stimulating Activity of PTH Fragments in LLC-PK1 Cells. A) Rat PTH(1-34) analog or amino-terminal rPTH fragments were tested for cAMP-stimulating activity in an LLC-P1-derived cell line (HKRK-B7) stably

transfected with the human PTH-1 receptor. Cells were treated with peptide at the indicated doses for 60 min at 22°C. Intracellular cAMP was measured by RIA, as described in Experimental Procedures. Shown are combined data (mean  $\pm$  s.e.m.) from 3 separate experiments, each performed in duplicate. B) HKRK-B7 cells, or untransfected LLC-PK1 were treated with rPTH(1-34) or rPTH(1-14), and intracellular cAMP was measured. Shown are data (mean  $\pm$  s.e.m.) from a single representative experiment performed in duplicate.

Figure 3. Alanine-scan of PTH(1-14). HKRK-B7 cells were treated with 100 mM of one of 14 different rPTH(1-14) analogs, each having a different alanine-substitution at the indicated amino acid position. The resulting cAMP levels were determined as described in Experimental Procedures. Shown are the combined data (mean  $\pm$  s.e.m.) from three separate experiments, each performed in duplicate. The mean (mean  $\pm$  s.e.m.) basal cAMP levels observed in the three experiments was  $2.1 \pm 0.1$  pmole/well, and the maximum response to rPTH(1-34) at 0.1 mM was  $254 \pm 16$  pmole/well.

Figure 4. PTH Responses of Intact and Truncated PTH-1 Receptors in COS-7 Cells. Shown at the top are schematics of the intact (A) and truncated (B) rat PTH-1 receptors used for transient transfection of COS-7 cells, and subsequent cAMP response assays. The conserved extracellular cysteine residues are depicted as open circles and numbered according to sequence position, and the nine amino acids of the epitope tag (HA) in rWT-HA are shaded. The ticks at residue 26 and 181 indicate the endpoints of the deletion in r $\delta$ Nt. Based on the predicted signal peptide cleavage site at Ala-22, residues 23-25 in r $\delta$ Nt are joined to residue 182. The cAMP responses of COS-7 cells expressing the intact receptor (C) and r $\delta$ Nt (D) to rPTH(1-34) (J) or rPTH(1-14) (C) are also shown. The graphs show combined data (mean  $\pm$  s.e.m.) from five separate experiments, each performed in duplicate.

Figure 5. Alanine-scan of PTH(1-14) with Intact and Truncated PTH Receptors. COS-7 cells transiently transfected with rWT-HA (A) or rCE I-G (B) were treated with 100 mM of native rat PTH(1-14) or 100 mM of a rPTH(1-14)

analog containing a single alanine substitution for 1h at 21 °C, and the resulting intracellular cAMP levels were measured by RIA. The amino acid substitutions are indicated on the axis labels. Peptides were tested in duplicate, and a single experiment representative of three others is shown.

5                   Figure 6. Specificity of the Truncated Ligand and PTH Receptor. COS-7 cells transiently transfected with either rWT-HA (A), rδNt (B), or the intact rat secretin receptor (C), were treated with the indicated peptides for 60 min at 22 °C, and the resulting intracellular cAMP levels were quantified by RIA. Concentration of peptides present during the incubations were: rPTH(1-34), 0.1 mM; rPTH(1-14) 100 mM; secretin(1-27), 1 mM and secretin(1-13), 100 mM. Shown are data (mean ± s.e.m.) from one experiment performed in duplicate, and this was repeated twice more with equivalent results.

10                   Figure 7. Antagonist Properties of PTHrP(7-34) with PTH(1-14). COS-7 cells transfected with rWT-HA (A) or rδNt (B) were treated with the antagonist [Leu11,D-Trp12]hPTHrP(7-34)NH<sub>2</sub> (or buffer alone), for 5 min at 22 °C, followed by 10 ml of either rPTH(1-34) or rPTH(1-14) agonist peptide. Incubations were continued for 30 min at 21 °C and the resulting cAMP levels were measured by RIA, as described in Experimental Procedures. The final concentration of antagonist peptide present during the incubation was 10 mM. Shown are data from a single experiment performed in triplicate. A repeat of the same experiment yielded equivalent results.

### ***Detailed Description of the Preferred Embodiments***

25                   The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding the rδNt receptor polypeptide, a novel, mutant PTH1R receptor polypeptide, having the amino acid sequence shown in Figure 1 (SEQ ID NO:2), which was determined by sequencing a cloned cDNA. The rδNt protein of the present invention shares sequence homology with previously identified non-mutant PTH1R and PTH2R sequences. The nucleotide sequence shown in Figure

1 (SEQ ID NO:1) was obtained by sequencing a cDNA clone (r $\delta$ Nt), which was deposited on \_\_\_\_\_ at the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209, and given accession number \_\_\_\_\_.

***1. The r $\delta$ Nt Receptor***

***a) Nucleic Acid Molecules***

Unless otherwise indicated, all nucleotide sequences determined by sequencing a DNA molecule herein were determined by manual sequencing, and all amino acid sequences of polypeptides encoded by DNA molecules determined herein were predicted by translation of a DNA sequence determined as above. Therefore, as is known in the art for any DNA sequence determined by this approach, any nucleotide sequence determined herein may contain some errors. Nucleotide sequences determined by manual sequencing are typically at least about 95% to at least about 99.9% identical to the actual nucleotide sequence of the sequenced DNA molecule. As is also known in the art, a single insertion or deletion in a determined nucleotide sequence compared to the actual sequence will cause a frame shift in translation of the nucleotide sequence such that the predicted amino acid sequence encoded by a determined nucleotide sequence will be completely different from the amino acid sequence actually encoded by the sequenced DNA molecule, beginning at the point of such an insertion or deletion.

Using the information provided herein, such as the nucleotide sequence in Figures 1, a nucleic acid molecule of the present invention encoding a r $\delta$ Nt polypeptide, respectively, may be obtained using standard techniques. Cloning and screening procedures are known for the isolation of the wild-type PTH1R sequence, such as those for cloning cDNAs using mRNA as starting material. Subsequent to cloning the wild-type receptor, the appropriate deletion in the sequence may be made as described herein. Illustrative of the invention, the nucleic acid molecule described in Figure 1 (SEQ ID NO:1) was obtained by using standard restriction enzyme digestion and cloning techniques in the art. The determined nucleotide sequence of the r $\delta$ Nt cDNA of Figure 1 (SEQ ID NO:1)

contains an open reading frame encoding a protein of about 435 amino acid residues, with a predicted leader sequence of about 22 amino acid residues. The amino acid sequence of the predicted mature rδNt receptor is shown in Figure 1 from amino acid residue about 23 to residue about 435. The rδNt protein shown in Figure 1 (SEQ ID NO:2) is about 84% identical to the rat PTH1 receptor.

As indicated, the present invention also provides the mature form(s) of the rδNt receptor of the present invention. According to the signal hypothesis, proteins secreted by mammalian cells have a signal or secretory leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Most mammalian cells and even insect cells cleave secreted proteins with the same specificity. However, in some cases, cleavage of a secreted protein is not entirely uniform, which results in two or more mature species on the protein. Further, it has long been known that the cleavage specificity of a secreted protein is ultimately determined by the primary structure of the complete protein, that is, it is inherent in the amino acid sequence of the polypeptide. Therefore, the present invention provides a nucleotide sequence encoding the mature rδNt polypeptides having the amino acid sequence encoded by the cDNA clone contained in the host identified as ATCC Deposit No. \_\_\_\_ and as shown in Figure 1 (SEQ ID NO:2). By the mature rδNt protein having the amino acid sequence encoded by the cDNA clone contained in the host identified as ATCC Deposit \_\_\_\_ is meant the mature form(s) of the rδNt receptor produced by expression in a mammalian cell (e.g., COS cells, as described below) of the complete open reading frame encoded by the DNA sequence of the clone contained in the vector in the deposited host. As indicated below, the mature rδNt receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. \_\_\_\_ may or may not differ from the predicted "mature" rδNt protein shown in Figure 1 (amino acids from about 23 to about 435) depending on the accuracy of the predicted cleavage.

Methods for predicting whether a protein has a secretory leader as well as the cleavage point for that leader sequence are available. For instance, the

methods of McGeoch (*Virus Res.* 3:271-286 (1985)) and von Heinje (*Nucleic Acids Res.* 14:4683-4690 (1986)) can be used. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%. von Heinje, *supra*. However, the two methods do not always produce the same predicted cleavage point(s) for a given protein. A computational method may be found in the computer program "PSORT" (K. Nakai and M. Kanehisa, *Genomics* 14:897-911 (1992)), which is an expert system for predicting the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated.

In the present case, the predicted amino acid sequence of the complete rδNt polypeptide of the present invention was analyzed for structural properties by comparison to the rat rδNt sequence. This analysis provided predicted a cleavage site between amino acids 22 and 23 in Figure 1 (SEQ ID NO:2). Thus, the leader sequence for the rδNt receptor protein is predicted to consist of amino acid residues 1-22 in Figure 1 (amino acids 1 to 22 in SEQ ID NO:2), while the predicted mature rδNt protein consists of residues 23-435 (amino acids 23 to 435 in SEQ ID NO:2).

As indicated, nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA or RNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

As one of ordinary skill would appreciate, however, due to the possibilities of sequencing errors, the rδNt receptor polypeptide encoded by the deposited cDNA comprises about 435 amino acids, but may be anywhere in the range of 425-435 amino acids; and the leader sequence of this protein is about 22 amino acids, but may be anywhere in the range of about 10 to about 30 amino acids.

As indicated, nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA or RNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Isolated nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) shown in Figure 1 (SEQ ID NO:1); DNA molecules comprising the coding sequence for the rδNt receptor shown in Figure 1 (SEQ ID NO:2); and DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode the rδNt receptor. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate such degenerate variants.

In another aspect, the invention provides isolated nucleic acid molecules encoding the rδNt polypeptide having an amino acid sequence encoded by the cDNA clone contained in the plasmid deposited as ATCC Deposit No. \_\_\_\_ on \_\_\_\_\_. Preferably, the nucleic acid molecule will encode the mature polypeptide encoded by the above-described deposited cDNA clone. In a further embodiment, a nucleic acid molecule is provided encoding the rδNt polypeptide or the rδNt polypeptide lacking the N-terminal methionine. The invention also provides an

isolated nucleic acid molecule having the nucleotide sequence shown in SEQ ID NO:1 or the nucleotide sequence of the rδNt cDNA contained in the above-described deposited clone, or a nucleic acid molecule having a sequence complementary to one of the above sequences. Such isolated molecules, particularly DNA molecules, are useful as probes for gene mapping, by *in situ* hybridization with chromosomes, and for detecting expression of the rδNt gene in human tissue, for instance, by Northern blot analysis.

The present invention is further directed to fragments of the isolated nucleic acid molecules described herein. By a fragment of an isolated nucleic acid molecule having the nucleotide sequence of the deposited cDNAs or the nucleotide sequence shown in Figure 1 (SEQ ID NO:1) is intended fragments at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length which are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments of about 50-1550 nt in length, and more preferably at fragments least about 600 nt in length are also useful according to the present invention as are fragments corresponding to most, if not all, of the nucleotide sequence of the deposited cDNAs or as shown in Figure 1 (SEQ ID NO:1). By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from the nucleotide sequence of the deposited cDNAs or the nucleotide sequence as shown in Figure 1 (SEQ ID NO:1).

Preferred nucleic acid fragments of the present invention include nucleic acid molecules encoding: a polypeptide comprising the rδNt receptor extracellular domain (predicted to constitute amino acid residues from about 23 to about 147 in Figure 1 (or amino acid residues from about 23 to about 147 in SEQ ID NO:2)); a polypeptide comprising the rδNt receptor transmembrane domain (predicted to constitute amino acid residues from about 148 to about 416 in Figure 1 (or amino acid residues from about 148 to about 416 in SEQ ID NO:2)); and a polypeptide comprising the rδNt receptor extracellular domain with all or part of the transmembrane domain deleted. As above with the leader sequence,

the amino acid residues constituting the rδNt receptor extracellular and transmembrane domains have been predicted. Thus, as one of ordinary skill would appreciate, the amino acid residues constituting these domains may vary slightly (e.g., by about 1 to about 15 amino acid residues) depending on the criteria used to define each domain.

Preferred nucleic acid fragments of the present invention also include nucleic acid molecules encoding epitope-bearing portions of the rδNt receptor protein. As one skilled in the art would know, a nucleic acid sequence may be used to predict the polypeptide sequence encoded therein. Such information may then be used to predict antigenic determinants in the polypeptide that may be related to the corresponding polynucleotide regions encoding the antigenic determinants identified by the analysis. Methods for predicting the antigenic determinants of a polypeptide are well known in the art.

Methods for determining other such epitope-bearing portions of the rδNt protein are described in detail below.

In another aspect, the invention provides an isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a portion of the polynucleotide in a nucleic acid molecule of the invention described above, for instance, the cDNA clones contained in ATCC Deposit Nos. \_\_\_\_\_ or \_\_\_\_\_. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150 mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 g/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers as discussed above and in more detail below.

By a portion of a polynucleotide of "at least 20 nt in length." for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the reference polynucleotide (e.g., the deposited cDNAs or the nucleotide sequence as shown in Figure 1 (SEQ ID NO:1).

5 Of course, a polynucleotide which hybridizes only to a poly A sequence (such as the 3' terminal poly(A) tract of the rδNt receptor cDNA shown in Figure 1 (SEQ ID NO:1), or to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of  
10 a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

As indicated, nucleic acid molecules of the present invention which encode a rδNt polypeptide may include, but are not limited to those encoding the amino acid sequence of the mature polypeptides, by themselves; the coding sequence for  
15 the mature polypeptides and additional sequences, such as those encoding the amino acid leader or secretory sequence, such as a pre-, or pro- or prepro- protein sequence; the coding sequence of the mature polypeptide, with or without the aforementioned additional coding sequences, together with additional, non-coding sequences, including for example, but not limited to introns and non-coding 5' and  
20 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription, mRNA processing, including splicing and polyadenylation signals, for example - ribosome binding and stability of mRNA; an additional coding sequence which codes for additional amino acids, such as those which provide additional functionalities. Thus, the sequence encoding the polypeptide may be  
25 fused to a marker sequence, such as a sequence encoding a peptide which facilitates purification of the fused polypeptide. In certain preferred embodiments of this aspect of the invention, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (Qiagen, Inc.), among others, many of which are commercially available. As described in Gentz *et al.*, *Proc.*  
30 *Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for

convenient purification of the fusion protein. The "HA" tag is another peptide useful for purification which corresponds to an epitope derived from the influenza hemagglutinin protein, which has been described by Wilson *et al.*, *Cell* 37: 767 (1984). As discussed below, other such fusion proteins include the rδNt receptor fused to Fc at the amino or C-terminus.

The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of the rδNt receptor. Variants may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. *Genes II*, Lewin, B., ed., John Wiley & Sons, New York (1985). Non-naturally occurring variants may be produced using art-known mutagenesis techniques.

Such variants include those produced by nucleotide substitutions, deletions or additions, which may involve one or more nucleotides. The variants may be altered in coding regions, non-coding regions, or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the rδNt receptor or portions thereof. Also especially preferred in this regard are conservative substitutions.

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 95%, 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the full-length rδNt polypeptide having the complete amino acid sequence in SEQ ID NO:2, including the predicted leader sequence; (b) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:2, but lacking the N-terminal methionine; (c) a nucleotide sequence encoding the mature rδNt receptor (full-length polypeptide with the leader removed) having the amino acid sequence at positions from about 23 to about 435 in SEQ ID NO:2; (d) a nucleotide sequence encoding the full-length rδNt polypeptide having the

complete amino acid sequence including the leader encoded by the cDNA clone contained in ATCC Deposit No. \_\_\_\_; (e) a nucleotide sequence encoding the mature rδNt receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97883; (f) a nucleotide sequence encoding the rδNt receptor extracellular domain; (g) a nucleotide sequence encoding the rδNt receptor transmembrane domain; (h) a nucleotide sequence encoding the rδNt receptor extracellular domain with all or part of the transmembrane domain deleted; and (i) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g) or (h).

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding a rδNt polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the rδNt receptor. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least 95%, 96%, 97%, 98% or 99% identical to, for instance, the nucleotide sequence shown in Figure 1 or to the nucleotides sequence of the deposited cDNA clones can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive,

Madison, WI 53711. Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

The present application is directed to nucleic acid molecules at least 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequence shown in Figure 1 (SEQ ID NO:1) or to the nucleic acid sequence of the deposited cDNAs, irrespective of whether they encode a polypeptide having rδNt receptor activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having rδNt receptor activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having rδNt receptor activity include, *inter alia*, (1) isolating the rδNt receptor gene or allelic variants thereof in a cDNA library; (2) *in situ* hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the rδNt receptor gene, as described in Verma *et al.*, *Human Chromosomes: A Manual of Basic Techniques*, Pergamon Press, New York (1988); and (3) Northern Blot analysis for detecting rδNt receptor mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequence shown in Figure 1 (SEQ ID NO:1) or to the nucleic acid sequence of the deposited cDNA which do, in fact, encode a polypeptide having rδNt receptor activity. By "a polypeptide having rδNt receptor activity" is intended polypeptides exhibiting activity similar, but not necessarily identical, to an activity of the rδNt receptor of

the invention, as measured in a particular biological assay. For example, rδNt receptor activity can be measured using competition binding experiments of labeled PTH or PTHrP to cells expressing the candidate rδNt polypeptide as described herein.

5 Any cell line expressing the rδNt receptor, or variants thereof, may be used to assay ligand binding and second messenger activation as described in Examples 3 and 4. Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 95%, 96%, 97%, 98%, or 99% identical to  
10 the nucleic acid sequence of the deposited cDNAs or the nucleic acid sequence shown in Figure 1 (SEQ ID NO:1) will encode a polypeptide "having rδNt receptor activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further  
15 recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having rδNt protein activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino  
20 acid).

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that proteins are surprisingly  
25 tolerant of amino acid substitutions.

#### ***b) Vectors and Host Cells***

The present invention also relates to vectors which include the isolated DNA molecules of the present invention, host cells which are genetically

engineered with the recombinant vectors, and the production of rδNt polypeptides or fragments thereof by recombinant techniques.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged *in vitro* using an appropriate packaging cell line and then transduced into host cells.

The DNA insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the *E. coli lac*, *trp* and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the constructs will preferably include a translation initiating at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase or neomycin resistance for eukaryotic cell culture and tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1

and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

5 Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods In Molecular Biology* (1986).

10 The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals, but also additional heterologous functional regions. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence in the host cell, during purification, or during subsequent handling and storage. Also, peptide moieties  
15 may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability and to facilitate purification, among others, are familiar and routine techniques in the art. A preferred fusion protein comprises a heterologous region from  
20 immunoglobulin that is useful to solubilize proteins. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is thoroughly advantageous for use in therapy and diagnosis and thus results, for  
25 example, in improved pharmacokinetic properties (EP-A 0232 262). On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified in the advantageous manner described. This is the case when Fc portion proves to be a hindrance to use in therapy and diagnosis, for example when the fusion protein is to be used as  
30 antigen for immunizations. In drug discovery, for example, human proteins, such

as. hIL5-receptor has been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett *et al.*, *Journal of Molecular Recognition*, Vol. 8:52-58 (1995) and K. Johanson *et al.*, *The Journal of Biological Chemistry*, Vol. 270, No. 16:9459-9471 (1995).

The rδNt receptor can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification. Polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

**c) *rδNt Polypeptides and Fragments***

The invention further provides an isolated rδNt polypeptide having the amino acid sequence encoded by the deposited cDNAs, or the amino acid sequence in Figure 1 (SEQ ID NO:2) or a peptide or polypeptide comprising a portion of the above polypeptides.

It will be recognized in the art that some amino acid sequences of the rδNt receptor can be varied without significant effect of the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on the protein which determine

activity. Thus, the invention further includes variations of the rδNt receptor which show substantial rδNt receptor activity or which include regions of rδNt protein such as the protein portions discussed below. Such mutants include deletions, insertions, inversions, repeats, and type substitutions. As indicated above, guidance concerning which amino acid changes are likely to be phenotypically silent can be found in Bowie, J.U., *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990).

Thus, the fragment, derivative or analog of the polypeptide of Figure 1 (SEQ ID NO:2) or that encoded by the deposited cDNAs, may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide, such as an IgG Fc fusion region peptide or leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

Of particular interest are substitutions of charged amino acids with another charged amino acid and with neutral or negatively charged amino acids. The latter results in proteins with reduced positive charge to improve the characteristics of the rδNt protein. The prevention of aggregation is highly desirable. Aggregation of proteins not only results in a loss of activity but can also be problematic when preparing pharmaceutical formulations, because they can be immunogenic. (Pinckard *et al.*, *Clin Exp. Immunol.* 2:331-340 (1967); Robbins *et al.*, *Diabetes*

36:838-845 (1987); Cleland *et al. Crit. Rev. Therapeutic Drug Carrier Systems* 10:307-377 (1993)).

The replacement of amino acids can also change the selectivity of binding to cell surface receptors. Ostade *et al. Nature* 361:266-268 (1993) describes certain mutations resulting in selective binding of TNF- $\alpha$  to only one of the two known types of TNF receptors. Thus, the r $\delta$ Nt receptor of the present invention may include one or more amino acid substitutions, deletions or additions, either from natural mutations or human manipulation.

As indicated, changes are preferably of a minor nature, such as conservative amino acid substitutions that do not significantly affect the folding or activity of the protein (see Table 1).

TABLE 1. Conservative Amino Acid Substitutions.

Aromatic	Phenylalanine Tryptophan Tyrosine
Hydrophobic	Leucine Isoleucine Valine
Polar	Glutamine Asparagine
Basic	Arginine Lysine Histidine
Acidic	Aspartic Acid Glutamic Acid
Small	Alanine Serine Threonine Methionine Glycine

Amino acids in the r $\delta$ Nt protein of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations

at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as receptor binding or *in vitro* proliferative activity. Sites that are critical for ligand-receptor binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992) and de Vos *et al. Science* 255:306-312 (1992)).

The polypeptides of the present invention are preferably provided in an isolated form. By "isolated polypeptide" is intended a polypeptide removed from its native environment. Thus, a polypeptide produced and/or contained within a recombinant host cell is considered isolated for purposes of the present invention. Also intended as an "isolated polypeptide" are polypeptides that have been purified, partially or substantially, from a recombinant host cell. For example, a recombinantly produced version of the antimicrobial peptide polypeptide can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988).

The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of the rδNt receptor can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988).

The polypeptides of the present invention also include the polypeptide encoded by the deposited rδNt cDNA including the leader, the polypeptide encoded by the deposited the cDNA minus the leader (i.e., the mature protein), the polypeptide of Figure 1 (SEQ ID NO:2) including the leader, the polypeptide of Figure 1 (SEQ ID NO:2) minus the leader, the extracellular domain, the transmembrane domain, a polypeptide comprising amino acids about 1 to about 435 in SEQ ID NO:2, and a polypeptide comprising amino acids about 2 to about 435 in SEQ ID NO:2, as well as polypeptides which are at least 95% identical, still more preferably at least 96%, 97%, 98% or 99% identical to the polypeptides described above, and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a reference amino acid sequence of a rδNt polypeptide is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of the rδNt receptor. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence shown in Figure 1 (SEQ ID NO:2) to the amino acid sequence encoded by deposited cDNA clones can be determined conventionally using known computer programs such the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

## 2. *Agonists and Antagonists of the rδNt Receptor Activity*

Functional characterization of the biological properties of the rδNt receptor of the invention and derivatives thereof may be performed by bioassays that measure ligand-stimulated cAMP accumulation.

**A. Assay for the Detection of Cyclic AMP Accumulation in Cells  
Expressing rδNt Receptor After Exposure to Test Compounds**

Intracellular cAMP accumulation is measured as described previously (Abou-Samra *et al.*, *J. Biol. Chem.* 262:1129, 1986). Cells expressing the rδNt receptor grown in 24-well plates are rinsed with culture medium containing 0.1% BSA and 2 mM IBMX. The cells are then incubated with a test compound for 60 min. at 21 °C. The supernatant is removed and the cells immediately frozen by placing the whole plate in dry ice powder. Intracellular cAMP is extracted by thawing the cells in 1 ml of 50 mM HCl and analyzed by a specific radioimmunoassay using an anti-cAMP antibody (e.g., Sigma, St. Louis, Mo). A cAMP analog (2'-O-monosuccinyl-adenosine 3':5'-cyclic monophosphate tyrosyl methyl ester, obtained from Sigma) which is used a tracer for cAMP is iodinated by the chloramine T method. Free iodine is removed by adsorbing the iodinated cAMP analog onto a C18 Sep-pak cartridge (Waters, Milford, Mass.). After washing with dH<sub>2</sub>O, the iodinated cAMP analog is eluted from the Sep-pak Cartridge with 40% acetonitrille (ACN) and 0.1% trifluoroacetic acid (TFA). The iodinated cAMP analog is lyophilized, reconstituted in 1 ml 0.1% TFA, and injected into a C18 reverse phase HPLC column (Waters). The column is equilibrated with 10% ACN in 0.1% TFA, and eluted with gradient of 10-30% ACN in 0.1% TFA. This allows separation of the mono-iodinated cAMP analog from the non-iodinated cAMP analog. The tracer is stable for up to 4 months when stored at - 20 °C. The standard used for the assay, adenosine 3':5'-cyclic monophosphate, may be purchased from Sigma. Samples (1-10 82 l of HCl extracts) or standards (0.04-100 fmol/tube) are diluted in 50 mM Na-acetate (pH 5.5), and acetylated with 10 µl of mixture of triethylamine and acetic anhydride (2:1 vol:vol). After acetylation, cAMP antiserum (100 µl) is added from a stock

solution (1:4000) made in PBS (pH 7.4), 5 mM EDTA and 1% normal rabbit serum. The tracer is diluted in PBS (pH 7.4) with 0.1% BSA, and added (20,000 cpm/tube). The assay is incubated at 4°C. overnight. The bound tracer is precipitated by adding 100 µl of goat anti-rabbit antiserum (1:20 in PBS) and 1 ml of 7% polyethyleneglycol (MW 5000-6000), centrifuging at 2000 rpm for 30 min. at 4°C. The supernatant is removed and the bound radioactivity is counted in a gamma-counter (Micromedic). To compute the cAMP data, logit calculations are performed in Excel spreadsheets. Typically, the assay sensitivity is 0.1 fmol/tube, and the standard concentration that displaces 50% of tracer is 5 fmol/tube.

***B. Screening Compounds Utilizing an rδNt Receptor Binding Assay***

In addition to the cAMP accumulation assay described below, it is possible that compounds may be iodinated and used in a radioreceptor-based assay in rδNt transiently transfected COS cells. COS-7 cells are grown in 15 cm plates in DMEM, 10% heat-inactivated FBS, 10 mg/L gentamycin until 80-90% confluent. Twenty-four hours after transfection by the DEAE/Dextran method (Sambrook *et al.*, *supra*), with 1-2µg of plasmid DNA, the cells are trypsinized and replated in multiwell plastic dishes (16 or 35 mm diameter, Costar, Cambridge, Mass.) at a cell concentration of  $5 \times 10^4$  cells/cm<sup>2</sup>. Cell number increased only slightly after transfection. After continuing culture for another 48 h, radioreceptor assays are performed. The culture medium is replaced with buffer containing 50 mM Tris-HCL (pH 7.7), 100 mM NaCl, 2 mM CaCl<sub>2</sub>, 5 mM KCL, 0.5% heat-inactivated fetal bovine serum (GIBCO), and 5% heat-inactivated horse serum (KC Biological Inc., Lenexa, Kans.) immediately before studies are initiated. Unless otherwise indicated, studies are conducted with cells incubated in this buffer at 15°C. for 4 h with  $4 \times 10^5$  cpm/ml ( $9.6 \times 10^{-11}$  M) of <sup>125</sup>I-labeled [Ala<sup>1</sup>]PTH(1-14) amide or <sup>125</sup>I-labeled [Nle<sup>8</sup>]PTH(1-14).

**C. Screening for PTH-1 Receptor Antagonists and Agonists**

The rδNt receptor of the invention may be utilized in to screen for compounds that are agonistic or antagonistic to the PTH response using the cAMP accumulation assay. Cells expressing PTH-1 receptor on the cell surface are incubated with native PTH(1-84) for 5-60 minutes at 37°C., in the presence of 2 mM IBMX (3-isobutyl-1-methyl-xanthine, Sigma, St. Louis, MO). Cyclic AMP accumulation is measured by specific radio-immunoassay, as described above. A test compound that competes with native PTH(1-84) for binding to the rδNt receptor, and that inhibits the effect of native PTH(1-84) on cAMP accumulation, is considered a competitive antagonist. Such a compound would be useful for treating hypercalcemia.

Conversely, a test compound that does not compete with native PTH(1-84) for binding to the rδNt receptor, but which still prevents native PTH(1-84) activation of cAMP accumulation (presumably by blocking the receptor activation site) is considered a non-competitive antagonist. Such a compound would be useful for treating hypercalcemia.

A candidate compound that competes with native PTH(1-84) for binding to the rδNt receptor, and which stimulates cAMP accumulation in the presence or absence of native PTH(1-84) is a competitive agonist. A candidate compound that does not compete with native PTH(1-84) for binding to the rδNt receptor but which is still capable of stimulating cAMP accumulation in the presence or absence of native PTH(1-84), or which stimulates a higher cAMP accumulation than that observed with a PTH compound would be considered a non-competitive agonist.

Thus, in a further aspect, a screening method is provided for determining whether a candidate agonist or antagonist is capable of enhancing or inhibiting a cellular response to PTH or PTHrP. The method involves contacting cells which express the rδNt polypeptide with a candidate compound and the PTH or PTHrP ligand, assaying a cellular response, and comparing the cellular response to a standard cellular response, the standard being assayed when contact is made with

the ligand in absence of the candidate compound, whereby an increased cellular response over the standard indicates that the candidate compound is an agonist of the ligand/receptor signaling pathway and a decreased cellular response compared to the standard indicates that the candidate compound is an antagonist of the ligand/receptor signaling pathway. By "assaying a cellular response" is intended qualitatively or quantitatively measuring a cellular response to a candidate compound and/or PTH or PTHrP (e.g., cyclic AMP accumulation). By the invention, a cell expressing the rδNt polypeptide can be contacted with either an endogenous or exogenously administered PTH or PTHrP.

## Examples

### General Methods

**Peptides:** Peptides were prepared by the Biopolymer Synthesis Facility at Massachusetts General Hospital (Boston, MA) using solid-phase chemistry with Fmoc (N-(9-fluorenyl)methoxycarbonyl) protecting groups, and TFA-mediated cleavage and deprotection. All peptides were C-terminally amidated. The PTH(1-14) analogs were synthesized on a multiple peptide synthesizer (Advanced Chemtech Model 396 MBS) at 0.025 mM scale. The completed peptides were desalted by adsorption on a C18 cartridge (Sep-Pak) and then analyzed by reversed-phase C18-based HPLC, MALDI-mass spectrometry and amino acid analysis. The PTH(1-34) control peptide, [Nle<sup>8,21</sup>,Tyr<sup>34</sup>]rPTH-(1-34)NH<sub>2</sub>, and the PTHrP(7-34) antagonist peptide, [Leu<sup>11</sup>,D-Trp<sup>12</sup>]hPTHrP(7-34)NH<sub>2</sub>, were prepared on an Applied Biosystems Synthesizer (Model 431A) 0.1 mM scale, purified by reversed-phase C18-based HPLC and characterized as described above. Concentrated stock solutions of peptides, 10 mM for PTH(1-14) analogs and 0.3 mM for PTHrP(7-34) and PTH(1-34), were prepared in 10 mM acetic acid, quantified by acid hydrolysis and amino acid analysis and stored at -80°C.

**Cell Culture and DNA Transfection:** COS-7 and HKRK-B7 cells were cultured at 37°C in Dulbecco's modified Eagle's medium (DMEM) supplemented with fetal bovine serum (10%); penicillin G (20 units/ml), streptomycin sulfate (20

5  $\mu\text{g/ml}$ ) and amphotericin B ( $0.05 \mu\text{g/ml}$ ) in a humidified atmosphere containing 5%  $\text{CO}_2$ . Twenty-four to 16 hours prior to assay, cells in 24-well plates were shifted to a humidified incubator containing 5%  $\text{CO}_2$  that was set at  $33^\circ\text{C}$ . Stock solutions of EGTA/trypsin and antibiotics were from GIBCO; fetal bovine serum was from Hyclone Laboratories (Logan, UT). Derivation and characterization of the HKRK-B7 cell line by stable transfection of LLC-PK<sub>1</sub> cells with a pCDNA-1-based plasmid (In Vitrogen, San Diego, CA) encoding the hPTH-1 receptor was described previously (Takasu, H. and Bringhurst, F., *Endocrinology*, in press (1998)). These cells express PTH-1 receptors at a surface density of about  $1 \times 10^6$  PTH-binding sites per cell. The HKRK-B7 cells were used for functional assays 10 24 to 72 hours after the cell monolayer reached confluency.

For studies with the intact and truncated rat PTH-1 receptors, transient transfections of COS-7 cells were performed using DEAE-dextran as described previously (Bergwitz, C., *et al.*, *J. Biol. Chem.* 272:28861-28868)). The construction and initial characterization of the pCDNA-1 based plasmids encoding either the intact or truncated rat PTH-1 receptor has been described previously (Lee, C., *et al.*, *Endocrinology* 135(4):1488-1495)). The intact receptor (rWT-HA) contains a nine amino acid HA epitope tag in place of residues 93-101 of the extracellular domain; this epitope tag does not affect receptor function (Lee, C., *et al.*, *Endocrinology* 135(4):1488-1495)). The truncated rat PTH-1 receptor (r $\delta$ Nt) is deleted for exons E1 through exon G (residues 26 to 181). Assuming that signal peptidase cleavage occurs between Ala-22 and Tyr-23 (Nielsen, H., *et al.*, *Protein Engineering* 10:1-6 (1997)), r $\delta$ Nt is predicted to have for its N-terminus residues Dautzenberg, F., *et al.*, *Proc. Natl. Acad. Sci.* 95:4941-4946 (1998); Holtmann, M., *et al.*, *J. Biol. Chem.* 270:14394-14398 (1995); DeAlmeida, V. and Mayo, K., *Mol. Endo.* 12:750-765 (1998)) joined to Glu-182 (Figure 3B). COS-7 cells were transfected in 24-well plates when the cells were 85 to 95% of confluency, using 200 ng of plasmid DNA that was purified by cesium chloride/ethidium bromide gradient centrifugation for each well. Assays were conducted 72 to 96 hours after transfection. Under these conditions about

~20 % of the COS-7 cells become transfected and express about  $5 \times 10^6$  surface PTH receptors per cell (Bergwitz, C., *et al.*, *J. Biol. Chem.* 272:28861-28868)).

Intracellular cyclic AMP: Transfected COS-7 or HKRK-B7 cells were rinsed with 500 ml of binding buffer, (50 mM Tris-HCl, pH 7.7, 100 mM NaCl, 5 mM KCl, 2 mM  $\text{CaCl}_2$ , 5% heat-inactivated horse serum, 0.5% heat-inactivated fetal bovine serum) and 200 ml of IBMX buffer (DMEM containing 2 mM IBMX, 1 mg/ml bovine serum albumin, 35 mM Hepes-NaOH, pH 7.4) and 100 ml of binding buffer or binding buffer containing various amounts of peptide were added. The plates were incubated for 60 minutes at room temperature. The buffer was then withdrawn and the cells were frozen on dry ice, treated with 0.5 ml of 50 mM HCl, and refrozen. After the thawing, the lysate was diluted 30-fold in dH<sub>2</sub>O and an aliquot was analyzed for cAMP content by determined radioimmunoassay using unlabeled cAMP as a standard.

For cAMP inhibition assays, transfected COS-7 cells were rinsed once with 500 ml of binding buffer, and 200 ml of IBMX buffer and 100 ml of binding buffer or binding buffer containing the antagonist [Leu<sup>11</sup>,D-Trp<sup>12</sup>]hPTHrP(7-34)NH<sub>2</sub> (10 mM) were added. After a 5 minute incubation at room temperature, 10 ml of binding buffer containing PTH(1-14) or PTH(1-34) (agonist peptide) was added, and the incubation was continued for an additional 30 minutes. The cells were then lysed and intracellular cAMP levels were measured as described above.

### ***Example 1***

#### ***PTH(1-14) action in stable cells.***

Amino-terminal peptide fragments based on the rat PTH sequence and ranging in length from PTH(1-9) to PTH(1-15) were synthesized and tested for activity in an LLC-PK1-derived cell line called HKRK-B7 which stably expresses high levels ( $1 \times 10^6$  receptors/cell) of the cloned human PTH-1 receptor (Takasu, H. and Bringham, F., *Endocrinology*, in press (1998)). As shown in Figure 1A, the intact control peptide PTH(1-34) mediated a 50-fold increase in intracellular cAMP levels relative to the basal cAMP level, and the estimated EC<sub>50</sub> for this

response was ~2nM with PTH(1-13) and shorter fragments little or no increase in cAMP accumulation was observed (Figure 1A). However, two of the amino-terminal fragments, PTH(1-14) and PTH(1-15), stimulated cAMP formation to about 20-fold over the basal level, although the doses required for this activation were five to six orders of magnitude higher than the dose required for PTH(1-34). The response to these active peptides was dependent on the transfected PTH receptor, as parental LLC-PK1 cells, which do not express PTH receptors, but do express the related calcitonin receptor, were unresponsive to PTH(1-34) or PTH(1-14) (Figure 2B).

With the intact receptor, the potency of PTH(1-14) was about five orders of magnitude weaker than that of PTH(1-34). This reduced potency is not surprising, given that the PTH(1-14) peptide lacks important receptor-binding residues located in the PTH(15-34) region in the PTH(15-34) domain (Nussbaum, S.R., *et al.*, *J. Biol. Chem.* 255:10183-10187 (1980); Gardella, T.J., *et al.*, *Endocrinology* 132(5):2024-2030 (1993); Caulfield, M.P., *et al.*, *Endocrinology* 127:83-87 (1990); and Abou-Samra, A.-B., *et al.*, *Endocrinology* 125:2215-2217 (1989)). Consistent with this, unlabeled PTH(1-14) bound too weakly to permit detection in our standard competition binding assays which used radioiodinated rPTH(1-34) as a tracer ligand, nor could we detect direct binding of radiolabeled PTH(1-14) analog to the intact or truncated receptors used in this study (data not shown).

## Example 2

### Alanine Scanning of PTH(1-14)

To identify residues in the PTH(1-14) fragment that play a role in activating the adenylyl cyclase-signaling pathway, an alanine-scanning approach was employed. Thirteen different alanine-substituted rat PTH(1-14) analogs were synthesized and tested for the ability to stimulate cAMP formation in HKRK-B7 cells (Figure 2). The activity profile obtained with the monosubstituted analogs revealed that residues in the 1-9 region formed a relatively intolerant segment of

the peptide, whereas residues in the 10-14 region formed a comparatively tolerant segment. Thus, with the exception of Ser-3 and Ala-1 (which is the native amino-terminal residue of rat PTH) most alanine substitutions in the 1-9 region yielded peptides that were barely active or inactive. In contrast, each alanine substitution in the 10-14 region yielded peptides with activities comparable with that of native rat PTH(1-14). The activity of the alanine-3 substituted peptide correlates well with previous studies on PTH(1-34) analogs which showed that amino acids with small side chains are tolerated at this site (Cohen, F.E., *et al.*, *J. Biol. Chem.* 266:1997-2004 (1991)).

### Example 3

#### *Potency of the r $\delta$ Nt Receptor with Small Ligands*

Experiments utilized COS-7 cells transfected with either the intact rat PTH-1 receptor (rWT-HA, Fig. 4A) or a truncated rat PTH-1 receptor with most of the amino-terminal extracellular domain deleted (r $\delta$ Nt, Fig. 4B). In COS-7 cells expressing rWT-HA, PTH(1-34) and PTH(1-14) mediated cAMP responses that were similar to the responses seen in HKRK-B7 cells: PTH(1-14) stimulated a 15-fold in cAMP formation, but with a potency that was four to five orders of magnitude weaker than that of PTH(1-34) (Figure 3C). Both peptides also stimulated cAMP formation in cells transfected with r $\delta$ Nt, but the potency of PTH(1-14) was only two orders of magnitude weaker than that of PTH(1-34) with this truncated receptor (Figure 3 D). This change in the relative potency of the two ligands could be accounted for by a 100-fold decrease in the potency which PTH(1-34) exhibited with r $\delta$ Nt, as compared to its potency with rWT-HA, rather than a shift in the potency of PTH(1-14), which was equipotent with the two receptors (compare panels C and D of Figure 4).

Although there was no direct measurement of receptor expression in this study, the 100-fold reduction in potency that PTH(1-34) exhibited with r $\delta$ Nt, as compared to rWT-HA, is not likely to be due to a reduction in surface expression of the truncated receptor, since PTH(1-14) exhibited equivalent activity with r $\delta$ Nt

and rWT-HA (Figure 4, C and D). This suggests that the two receptors are expressed at approximately equal levels. The reduced activity of PTH(1-34) with rδNt therefore most likely reflects a loss of important binding interactions that normally occur between the (Jüppner, H., *et al.*, *Science* 254:1024-1026 (1991),  
5 Guo, J., *et al.*, *Endocrinology* 136:3884-3891 (1995); Hruska, K.A., *J. Clin. Invest.* 79:230-239 (1987); Donahue, H.J., *et al.*, *J. Biol. Chem.* 263:13522-13527 (1988); Kolakowski, L.F., *GCRDb: A G-protein-coupled receptor Database Receptors and Channels* 2:1-7 (1994); Jüppner, H., *et al.*, *Endocrinology* 134:879-884 (1994); Lee, C., *et al.*, *Mol. Endo.* 9:1269-1278  
10 (1995); Turner, P., *et al.*, Single Mutations Allow the PTH-2 Receptor to Respond to PTHrP *J. Bone Min. Res.* 12, Supplement 1, Abstract #121 (1997); Dautzenberg, F., *et al.*, *Proc. Natl. Acad. Sci.* 95:4941-4946 (1998), Holtmann, M., *et al.*, *J. Biol. Chem.* 270:14394-14398 (1995); DeAlmeida, V., *et al.*, *Mol. Endo.* 12:750-765 (1998); Stroop, S., *et al.*, *Biochem.* 34:1050-1057 (1994);  
15 Zhou, A., *et al.*, *Proc. Natl. Acad. Sci. USA* 94:3644-3649 (1997); Bisello, A., *et al.*, *J. Biol. Chem.* 273:22498-22505 (1998); Bergwitz, C., *et al.*, *J. Biol. Chem.* 271:26469-26472 (1996); Gardella, T.J., *et al.*, *Endocrinology* 135:1186-1194 (1994); Mannstadt, M., *et al.*, *J. Biol. Chem.* 273:16890-16896 (1998); Takasu, H. and Bringham, F., *Endocrinology*, in press (1998); Bergwitz, C., *et al.*, *J. Biol. Chem.* 272:28861-28868 (1997); Lee, C., *et al.*, *Endocrinology* 135(4):1488-1495 (1994)) domain of the ligand and the amino-terminal domain of the receptor (Jüppner, H., *et al.*, *Endocrinology* 134:879-884 (1994); Bergwitz, C., *et al.*, *J. Biol. Chem.* 271:16469-26472 (1996); Mannstadt, M., *et al.*, *J. Biol. Chem.* 273:16890-16896 (1998)). The lack of these same binding interactions  
20 could also explain the inability of PTHrP(7-34) to function as an antagonist with rδNt ( see Figure 7B).

#### **Example 4:**

***The rδNt Receptor Interacts with the same PTH(1-14)  
Functional Residues as Intact PTH-1 receptor***

Experiments were designed to test whether the PTH(1-14) residues that are required for function with the truncated receptor differ from those required for function with the intact receptor. Using the alanine-scanning set of PTH(1-14) analogs, experiments tested cAMP-stimulating activity for the two rat PTH-1  
5 receptors in COS-7 cells. As shown in Figures 5A and 5B, the activity profiles obtained with rδNt mirrored that obtained with rWT-HA, since Ser-3 and the 10-14 region of the peptide were tolerant of mutation, whereas residues 2 and 4-9 were intolerant (Figures 4A and B). Therefore, the same set of functional residues in PTH(1-14) that are required for interaction with the intact PTH-1 receptor are  
10 also required for interaction with the core domain of the receptor.

### ***Example 5***

#### ***Specificity of Truncated Ligands and PTH-receptors***

In order to test whether PTH(1-14) and rδNt retained the appropriate recognition specificity for the corresponding parent ligand, cross-reactivity  
15 experiments were done using secretin ligands and the cloned rat secretin receptor. COS-7 cells transfected with the secretin receptor exhibited a 50-fold increase in cAMP levels in response to secretin(1-27) (1 mM), but did not respond to either PTH(1-34) (1 mM) or PTH(1-14) (100 mM) (Figure 5C). Cells expressing rδNt responded to PTH(1-34) and PTH(1-14) but not to secretin(1-27) (1 mM) or  
20 secretin(1-13) (1 mM) (Figure 5B). Thus, the recognition specificity of PTH(1-14) and rδNt appear to replicate that of the intact parent molecules. No evidence for relaxed specificity was detected in these studies (Figure 6). It is also worth noting that PTH(1-14) did not activate the endogenous calcitonin receptors expressed in LLC-PK1 cells (Figure 2B).

### ***Example 6***

#### ***rδNt Stimulation is Not Affected by the Inhibitor***

#### ***[Leu11,D-Trp12]hPTHrP(7-34)NH<sub>2</sub>***

In order to determine whether [Leu11,D-Trp12]hPTHrP(7-34)NH<sub>2</sub>, a potent competitive antagonist of PTH(1-34) action (Nutt, R.F., *et al.*, *Endocrinology* 127:491-493 (1990)), could block the ability of PTH(1-14) to stimulate cAMP formation in COS-7 cells expressing either rWT-HA or rδNt (Figure 6). With rWT-HA the inhibitor peptide reduced the efficacy of both PTH(1-14) and PTH(1-34) by as much as 70 % as compared to the responses elicited by these agonists in the absence of inhibitor (Figure 6A). In contrast, PTHrP(7-34) had little or no effect on the ability of PTH(1-34) or PTH(1-14) to stimulate cAMP production in cells expressing rδNt.

The ability of PTHrP(7-34) to antagonize PTH(1-14) action on the intact receptor (Figure 6A) suggests that the receptor sites occupied by these two ligands overlap. This overlap could involve ligand residues 7-14 and some portion in the core region of the receptor. Any binding interactions that may occur between figure residues (7-34) and the core region of the receptor are, however, too weak to enable effective antagonism in the absence of the amino-terminal extracellular receptor domain.

These findings establish that a much smaller region of PTH(1-34) than heretofore appreciated can stimulate receptor activation, and that the amino-terminal portion of PTH the hormone interacts with the core region of the receptor containing the seven transmembrane helices and connecting loops, as previously hypothesized for intact PTH ligands and receptors (Lee, C., *et al.*, *Mol. Endo.* 9:1269-1278 (1995); Bisello, A., *et al.*, *J. Biol. Chem.* 273:22498-22505 (1998); Bergwitz, C., *et al.*, *J. Biol. Chem.* 271:26469-26472 (1996); Gardella, T.J., *et al.*, *Endocrinology* 135:1186-1194 (1994); Bergwitz, C., *et al.*, *J. Biol. Chem.* 272:28861-28868; Gardella, T., *et al.*, *J. Biol. Chem.* 271:12820-12825 (1996)). Furthermore, this component of the interaction is sufficient for receptor signaling. The hypothesis that the 15-34 region of PTH binds to the amino-terminal extracellular domain of the receptor does not exclude the possibility that this domain, which by itself does not stimulate cAMP formation (data not shown), also provides some binding energy by interacting with the core region of the

receptor. In fact, the ~100-fold greater potency that PTH(1-34) exhibits with rδNt, in comparison to the potency of PTH(1-14) with this receptor (Figure 4D), might well be due to such interactions. However, we can not exclude the alternative possibility that the 15-34 domain enhances the intrinsic signaling activity of the (1-14) segment, for example, by stabilizing a favorable secondary structure in the amino-terminal portion of the ligand. More specific information on the receptor recognition sites utilized by PTH and the structure of the receptor-bound ligand are required to distinguish between such possibilities.

Some recognition determinants have been identified in the amino-terminal extracellular domain, the extracellular loops and the transmembrane helices of the B family of receptors (Turner, P., *et al.*, Single Mutations Allow the PTH-2 Receptor to Respond to PTHrP *J. Bone Min. Res.* 12, Supplement 1, Abstract #121 (1997), Dautzenberg, F., *et al.*, *Proc. Natl. Acad. Sci.* 95:4941-4946 (1998), Holtmann, M., *et al.*, *J. Biol. Chem.* 270:14394-14398 (1995), Gardella, T.J., *et al.*, *Endocrinology* 135:1186-1194 (1994); Bergwitz, C., *et al.*, *J. Biol. Chem.* 272:28861-28868; Turner, P.R., *et al.*, *J. Biol. Chem.* 271(16):9205-9208 (1996)).

One distinguishing feature of the family B receptors is the amino-terminal extracellular domain, which is relatively large and contains a number of conserved residues, including six cysteines. It is thus intriguing that this domain of the PTH-1 receptor is not essential for ligand-dependent signal transduction, as evidenced by the results with the rδNt receptor.

Several other reports on other family B receptors provide additional evidence to suggest that the amino-terminal extracellular domains of these receptors may not be essential for functional expression. Large amino-terminal deletions in the calcitonin receptor (Unson, C., *et al.*, *J. Biol. Chem.* 270:27720-27727 (1995)) and in the growth hormone-releasing factor receptor (DeAlmedia, V. and Mayo, K., *Mol. Endo.* 12:750-765 (1998)) were compatible with expression, as assessed by immunologic methods, and a glucagon receptor lacking the amino-terminal domain and containing an activating mutation in helix 2 (HR-

178) exhibited constitutive cAMP-signaling activity (Hjorth, S., *et al.*, *Mol. Endo.* 12:78-86 (1998)). In these studies, however, evidence that the truncated receptor could interact with ligand, as we have found for the PTH-1 receptor, was not reported. In a separate study on the lutropin receptor, a group A receptor that  
5 binds the large glyco hormone human choriogonadotropin, it was observed that a deletion of the large amino-terminal extracellular domain yielded a receptor that could mediate a cAMP response to high doses of hCG (Ji, I.H. and Ji, T.H., *J. Biol. Chem.* 266(20):13076-13079 (1991)).

That the activity of PTH(1-14) was not affected by the deletion of the  
10 amino-terminal receptor domain suggests that the peptide interacts predominantly with the core region of the receptor. This conclusion is supported by the alanine-scanning experiments performed on PTH(1-14), in which the profile of tolerant and intolerant residues observed with rδNt was nearly the same as that obtained with the intact receptor (Figure 4). With each receptor, residues in the 10-14  
15 region of the ligand formed a tolerant segment, while residues in the 1-9 region, excluding 1 and 3, formed an intolerant segment. This pattern of critical and non-critical residues observed in the context of the PTH(1-14) fragment closely matches the patterns found previously in studies on longer-length PTH analogs (Cohen, F.E., *et al.*, *J. Biol. Chem.* 266:1997-2004 (1991); Gombert, F., *et al.*,  
20 in "Peptides: Chemistry, Structure and Biology Proceedings of the 14<sup>th</sup> American Peptide Symposium June 18-23, Kaumaya, P. and Hodges, R., eds., pp. 661-662, Mayflower Scientific Limited, Kingswinford, UK (1996); Gardella, T.J., *et al.*, *J. Biol. Chem.* 266:13141-13146 (1991)).

***What Is Claimed Is:***

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

5 (a) a nucleotide sequence encoding the rδNt receptor having the complete amino acid sequence at positions from about 1 to about 435 in SEQ ID NO:2;

(b) a nucleotide sequence encoding the rδNt receptor having the amino acid sequence at positions from about 2 to about 435 in SEQ ID NO:2;

10 (c) a nucleotide sequence encoding the mature rδNt receptor having the amino acid sequence at positions from about 23 to about 435 in SEQ ID NO:2;

(d) a nucleotide sequence encoding the rδNt receptor having the complete amino acid sequence encoded by the cDNA clone contained in  
15 ATCC Deposit No. \_\_\_\_;

(e) a nucleotide sequence encoding the mature rδNt receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. \_\_\_\_;

20 (f) a nucleotide sequence encoding the rδNt extracellular domain;

(g) a nucleotide sequence encoding the rδNt transmembrane domain;

(h) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f) or (g).

25 2. The nucleic acid molecule of claim 1 wherein said polynucleotide has the complete nucleotide sequence in Figure 1 (SEQ ID NO:1).

3. The nucleic acid molecule of claim 1 wherein said polynucleotide has the nucleotide sequence in Figure 1 (SEQ ID NO:1) encoding the rδNt receptor having the complete amino acid sequence in Figure 1 (SEQ ID NO:2).

5 4. The nucleic acid molecule of claim 1 wherein said polynucleotide has the nucleotide sequence in Figure 1 (SEQ ID NO:1) encoding the mature rδNt receptor having the amino acid sequence in Figure 1 (SEQ ID NO:2).

5. The nucleic acid molecule of claim 1 wherein said polynucleotide has the complete nucleotide sequence of the cDNA clone contained in ATCC Deposit No. \_\_\_\_.

10 6. The nucleic acid molecule of claim 1 wherein said polynucleotide has the nucleotide sequence encoding the rδNt receptor having the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. \_\_\_\_.

15 7. The nucleic acid molecule of claim 1 wherein said polynucleotide has the nucleotide sequence encoding the mature rδNt receptor having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. \_\_\_\_.

20 8. An isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a polynucleotide having a nucleotide sequence identical to a nucleotide sequence in (a), (b), (c), (d), (e), (f) or (g) of claim 1 wherein said polynucleotide which hybridizes does not hybridize under stringent hybridization conditions to a polynucleotide having a nucleotide sequence consisting of only A residues or of only T residues.

9. An isolated nucleic acid molecule comprising a polynucleotide which encodes the amino acid sequence of an epitope-bearing portion of a rδNt receptor having an amino acid sequence in (a), (b), (c), (d), (e), (f) or (g) of claim 1.

5 10. The isolated nucleic acid molecule of claim 1, which encodes the rδNt receptor extracellular domain.

11. The isolated nucleic acid molecule of claim 1, which encodes the rδNt receptor transmembrane domain.

10 12. A method for making a recombinant vector comprising inserting an isolated nucleic acid molecule of claim 1 into a vector.

13. A recombinant vector produced by the method of claim 12.

14. A method of making a recombinant host cell comprising introducing the recombinant vector of claim 13 into a host cell.

15. A recombinant host cell produced by the method of claim 14.

15 16. A recombinant method for producing a rδNt polypeptide, comprising culturing the recombinant host cell of claim 15 under conditions such that said polypeptide is expressed and recovering said polypeptide.

17. An isolated rδNt polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

20 (a) the amino acid sequence of the rδNt polypeptide having the complete amino acid sequence at positions from about 1 to about 435 in SEQ ID NO:2;

(b) the amino acid sequence of the rδNt polypeptide having the amino acid sequence at positions from about 2 to about 435 in SEQ ID NO:2;

(c) the amino acid sequence of the mature rδNt polypeptide having the amino acid sequence at positions from about 23 to about 435 in SEQ ID NO:2;

(d) the amino acid sequence of the rδNt polypeptide having the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. \_\_\_\_; and

(e) the amino acid sequence of the mature rδNt polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. \_\_\_\_.

18. An isolated antibody that binds specifically to a rδNt receptor polypeptide of claim 17.

19. A method of screening for agonists or antagonists of PTH receptor activity comprising:

- (a) contacting a test compound to cells expressing a PTH receptor; and
- (b) measuring the biological response of said cells.

1/8

ATGGGGGCGCGCCGGATCGCACCCAGCCTGGCGCTCCTACTCTGCTGCCAGTGCTCAGC  
a M G A A R I A P S L A L L L C C P V L S -

TCCGCATATGCGCTGGAGGTATTTGACCGCCTAGGCATGATCTACACCGTGGGATACTCC  
a S A Y A L E V F D R L G M I Y T V G Y S -

ATGTCTCTCGCCTCCCTCACGGTGGCTGTGCTCATCCTGGCCTATTTTAGGCGGCTGCAC  
a M S L A S L T V A V L I L A Y F R R L H -

TGCACGCGCAACTACATCCACATGCACATGTTTCCTGTCGTTTATGCTGCGCGCCGCGAGC  
a C T R N Y I H M H M F L S F M L R A A S -

ATCTTCGTGAAGGACGCTGTGCTCTACTCTGGCTTCACGCTGGATGAGGCCGAGCGCCTC  
a I F V K D A V L Y S G F T L D E A E R L -

ACAGAGGAAGAGTTGCACATCATCGCGCAGGTGCCACCTCCGCCGGCCGCTGCCGCCGTA  
a T E E E L H I I A Q V P P P P A A A A V -

GGCTACGCTGGCTGCCGCGTGGCGGTGACCTTCTTCCTCTACTTCCTGGCTACCAACTAC  
a G Y A G C R V A V T F F L Y F L A T N Y -

TACTGGATCCTGGTGGAGGGGCTGTACTTGCACAGCCTCATCTTCATGGCCTTTTTCTCA  
a Y W I L V E G L Y L H S L I F M A F F S -

GAGAAGAAGTACCTGTGGGGCTTCACCATCTTTGGCTGGGGTCTACCGGCTGTCTTCGTG  
a E K K Y L W G F T I F G W G L P A V F V -

GCTGTGTGGGTGCGGTGTCAGAGCAACCTTGGCCAACACTGGGTGCTGGGATCTGAGCTCC  
a A V W V G V R A T L A N T G C W D L S S -

GGGCACAAGAAGTGGATCATCCAGGTGCCCATCCTGGCATCTGTTGTGCTCAACTTCATC  
a G H K K W I I Q V P I L A S V V L N F I -

CTTTTTATCAACATCATCCGGGTGCTTGCCACTAAGCTTCGGGAGACCAATGCGGGCCGG  
a L F I N I I R V L A T K L R E T N A G R -

FIG. 1

SUBSTITUTE SHEET (RULE 26)

2/8

## FIG. 1- CONTINUED

TGTGACACCAGGCAGCAGTACCGGAAGCTGCTCAGGTCCACGTTGGTGCTCGTGCCGCTC  
a C D T R Q Q Y R K L L R S T L V L V P L -

TTTGGTGTGCACTACACCGTCTTCATGGCCTTGCCGTACACCGAGGTCTCAGGGACATTG  
a F G V H Y T V F M A L P Y T E V S G T L -

TGGCAGATCCAGATGCATTATGAGATGCTCTTCAACTCCTTCCAGGGATTTTTTGTGTC  
a W Q I Q M H Y E M L F N S F Q G F F V A -

ATCATATACTGTTTCTGCAATGGTGAGGTGCAGGCAGAGATTAGGAAGTCATGGAGCCGC  
a I I Y C F C N G E V Q A E I R K S W S R -

TGGACACTGGCGTTGGACTTCAAGCGCAAAGCACGAAGTGGGAGTAGCAGCTACAGCTAT  
a W T L A L D F K R K A R S G S S S Y S Y -

GGCCCAATGGTGTCTCACACGAGTGTGACCAATGTGGGCCCCCGTGCAGGACTCAGCCTC  
a G P M V S H T S V T N V G P R A G L S L -

CCCCTCAGCCCCCGCCTGCCTCCTGCCACTACCAATGGCCACTCCCAGCTGCCTGGCCAT  
a P L S P R L P P A T T N G H S Q L P G H -

GCCAAGCCAGGGGCTCCAGCCACTGAGACTGAAACCCTACCAGTCACTATGGCGGTTCCC  
a A K P G A P A T E T E T L P V T M A V P -

AAGGACGATGGATTCTTAACGGCTCCTGCTCAGGCCTGGATGAGGAGGCCTCCGGGTCT  
a K D D G F L N G S C S G L D E E A S G S -

GCGCGGCCGCTCCATTGTTGCAGGAAGGATGGGAAACAGTCATGTGACTGGGCACTAGG  
3579 -+-----+-----+-----+-----+-----+-----+----- 3638  
a A R P P P L L Q E G W E T V M \* -

3/8




FIG. 2B

FIG. 2A

4/8




FIG. 3

5/8




FIG. 4A




FIG. 4B




FIG. 4C

FIG. 4D

6/8




FIG. 5A




FIG. 5B

7/8




FIG. 6A




FIG. 6B




FIG. 6C

8/8




FIG. 7A




FIG. 7B

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13*bis*)

**A.** The indications made below relate to the microorganism referred to in the description on page 7, line 3; page 8, lines 19, 21 and 26; page 10, line 26; page 12, line 20; page 15, line 2; page 38, lines 15 and 18; page 39, lines 9, 13 and 16; page 41, lines 8 and 11.

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☐

Name of depository institution

American Type Culture Collection (ATCC)

Address of depository institution (*including postal code and country*)

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit    December 28, 1999

Accession Number    PTA-1136

**C. ADDITIONAL INDICATIONS** (*leave blank if not applicable*)

This information is continued on an additional sheet ☐

*Escherichia coli* plasmid TG-98

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (*if the indications are not for all designated States*)

**E. SEPARATE FURNISHING OF INDICATIONS** (*leave blank if not applicable*)

The indications listed below will be submitted to the international Bureau later (*specify the general nature of the indications, e.g., "Accession Number of Deposit"*)

For receiving Office use only

For International Bureau use only

☐ This sheet was received with the international application

☐ This sheet was received by the International Bureau on:

Authorized officer

Authorized officer

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on page <u>page 7, line 3; page 8, lines 19, 21 and 26; page 10, line 26; page 12, line 20; page 15, line 2; page 38, lines 15 and 18; page 39, lines 9, 13 and 16; page 41, lines 8 and 11.</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution American Type Culture Collection (ATCC)	
Address of depository institution (including postal code and country)  10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit     December 28, 1999	Accession Number     PTA-1136
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-98  In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

***Escherichia coli Plasmid TG-98*****AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**CANADA**

The applicant hereby requests that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the furnishing of a sample of deposited biological material referred to in the application only be effected to an independent expert nominated by the Commissioner of Patents.

**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent office or any person approved by the applicant in the individual case.

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Registration), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the National Board of Patents and Registration or any person approved by the applicant in the individual case.

**ICELAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Icelandic Patent Office), or has been finally decided upon by the Icelandic Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

***Escherichia coli Plasmid TG-98*****NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in Rule 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**NORWAY**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Norwegian Patent office or any person approved by the applicant in the individual case.

**SINGAPORE**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent office or any person approved by the applicant in the individual case.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution American Type Culture Collection (ATCC)	
Address of depository institution (including postal code and country)  10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit     December 28, 1999	Accession Number     PTA-1138
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-422  In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution  American Type Culture Collection (ATCC)	
Address of depository institution <i>(including postal code and country)</i>  10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit    December 28, 1999	Accession Number    PTA-1138
<b>C. ADDITIONAL INDICATIONS</b> <i>(leave blank if not applicable)</i> <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-422	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> <i>(if the indications are not for all designated States)</i>	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the international Bureau later <i>(specify the general nature of the indications, e.g., "Accession Number of Deposit")</i>	

<div style="text-align: center; border-bottom: 1px solid black; margin-bottom: 5px;">For receiving Office use only</div> <div style="padding: 5px;"> <input type="checkbox"/> This sheet was received with the international application         </div> <div style="padding: 5px; border-top: 1px solid black;">           Authorized officer         </div>	<div style="text-align: center; border-bottom: 1px solid black; margin-bottom: 5px;">For International Bureau use only</div> <div style="padding: 5px;"> <input type="checkbox"/> This sheet was received by the International Bureau on:         </div> <div style="padding: 5px; border-top: 1px solid black;">           Authorized officer         </div>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

***Escherichia coli* Plasmid TG-422****AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**CANADA**

The applicant hereby requests that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the furnishing of a sample of deposited biological material referred to in the application only be effected to an independent expert nominated by the Commissioner of Patents.

**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent office or any person approved by the applicant in the individual case.

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Registration), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the National Board of Patents and Registration or any person approved by the applicant in the individual case.

**ICELAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Icelandic Patent Office), or has been finally decided upon by the Icelandic Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

***Escherichia coli Plasmid TG-422*****NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in Rule 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**NORWAY**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Norwegian Patent office or any person approved by the applicant in the individual case.

**SINGAPORE**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent office or any person approved by the applicant in the individual case.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution American Type Culture Collection (ATCC)	
Address of depository institution <i>(including postal code and country)</i> 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit    December 28, 1999	Accession Number    PTA-1139
<b>C. ADDITIONAL INDICATIONS</b> <i>(leave blank if not applicable)</i> <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-433	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> <i>(if the indications are not for all designated States)</i>	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the international Bureau later <i>(specify the general nature of the indications, e.g., "Accession Number of Deposit")</i>	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution American Type Culture Collection (ATCC)	
Address of depository institution ( <i>including postal code and country</i> )  10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit     December 28, 1999	Accession Number     PTA-1139
<b>C. ADDITIONAL INDICATIONS</b> ( <i>leave blank if not applicable</i> ) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-433  In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> ( <i>if the indications are not for all designated States</i> )	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> ( <i>leave blank if not applicable</i> )	
The indications listed below will be submitted to the international Bureau later ( <i>specify the general nature of the indications, e.g., "Accession Number of Deposit"</i> )	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

***Escherichia coli* Plasmid TG-433****AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**CANADA**

The applicant hereby requests that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the furnishing of a sample of deposited biological material referred to in the application only be effected to an independent expert nominated by the Commissioner of Patents.

**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent office or any person approved by the applicant in the individual case.

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Registration), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the National Board of Patents and Registration or any person approved by the applicant in the individual case.

**ICELAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Icelandic Patent Office), or has been finally decided upon by the Icelandic Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

***Escherichia coli Plasmid TG-433*****NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in Rule 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**NORWAY**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Norwegian Patent office or any person approved by the applicant in the individual case.

**SINGAPORE**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent office or any person approved by the applicant in the individual case.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution American Type Culture Collection (ATCC)	
Address of depository institution ( <i>including postal code and country</i> ) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit    December 28, 1999	Accession Number    PTA-1140
<b>C. ADDITIONAL INDICATIONS</b> ( <i>leave blank if not applicable</i> ) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-454	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> ( <i>if the indications are not for all designated States</i> )	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> ( <i>leave blank if not applicable</i> )	
The indications listed below will be submitted to the international Bureau later ( <i>specify the general nature of the indications, e.g., "Accession Number of Deposit"</i> )	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution American Type Culture Collection (ATCC)	
Address of depository institution <i>(including postal code and country)</i>  10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit     December 28, 1999	Accession Number     PTA-1140
<b>C. ADDITIONAL INDICATIONS</b> <i>(leave blank if not applicable)</i> <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-454  In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> <i>(if the indications are not for all designated States)</i>	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the international Bureau later <i>(specify the general nature of the indications, e.g., "Accession Number of Deposit")</i>	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

***Escherichia coli Plasmid TG-454*****AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**CANADA**

The applicant hereby requests that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the furnishing of a sample of deposited biological material referred to in the application only be effected to an independent expert nominated by the Commissioner of Patents.

**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent office or any person approved by the applicant in the individual case.

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Registration), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the National Board of Patents and Registration or any person approved by the applicant in the individual case.

**ICELAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Icelandic Patent Office), or has been finally decided upon by the Icelandic Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

***Escherichia coli Plasmid TG-454*****NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in Rule 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**NORWAY**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Norwegian Patent office or any person approved by the applicant in the individual case.

**SINGAPORE**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent office or any person approved by the applicant in the individual case.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

**A.** The indications made below relate to the microorganism referred to in the description on page 12, line 20.

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☐

Name of depository institution

American Type Culture Collection (ATCC)

Address of depository institution *(including postal code and country)*

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit    December 28, 1999

Accession Number    PTA-1142

**C. ADDITIONAL INDICATIONS** *(leave blank if not applicable)*

This information is continued on an additional sheet ☐

*Escherichia coli* plasmid TG-462

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** *(if the indications are not for all designated States)*

**E. SEPARATE FURNISHING OF INDICATIONS** *(leave blank if not applicable)*

The indications listed below will be submitted to the international Bureau later *(specify the general nature of the indications, e.g., "Accession Number of Deposit")*

For receiving Office use only

☐ This sheet was received with the international application

Authorized officer

For International Bureau use only

☐ This sheet was received by the International Bureau on:

Authorized officer

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 12, line 20.

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☐

Name of depository institution

American Type Culture Collection (ATCC)

Address of depository institution (including postal code and country)

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit      December 28, 1999

Accession Number      PTA-1142

**C. ADDITIONAL INDICATIONS** (leave blank if not applicable)

This information is continued on an additional sheet ☐

*Escherichia coli* plasmid TG-462

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (if the indications are not for all designated States)

**E. SEPARATE FURNISHING OF INDICATIONS** (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")

For receiving Office use only

☐ This sheet was received with the international application

Authorized officer

For International Bureau use only

☐ This sheet was received by the International Bureau on:

Authorized officer

***Escherichia coli* Plasmid TG-462****AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**CANADA**

The applicant hereby requests that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the furnishing of a sample of deposited biological material referred to in the application only be effected to an independent expert nominated by the Commissioner of Patents.

**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent office or any person approved by the applicant in the individual case.

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Registration), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the National Board of Patents and Registration or any person approved by the applicant in the individual case.

**ICELAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Icelandic Patent Office), or has been finally decided upon by the Icelandic Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

***Escherichia coli* Plasmid TG-462****NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in Rule 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**NORWAY**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Norwegian Patent office or any person approved by the applicant in the individual case.

**SINGAPORE**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent office or any person approved by the applicant in the individual case.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution  American Type Culture Collection (ATCC)	
Address of depository institution <i>(including postal code and country)</i>  10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit    December 30, 1999	Accession Number    PTA-1141
<b>C. ADDITIONAL INDICATIONS</b> <i>(leave blank if not applicable)</i> <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-449	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> <i>(if the indications are not for all designated States)</i>	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the international Bureau later <i>(specify the general nature of the indications, e.g., "Accession Number of Deposit")</i>	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution American Type Culture Collection (ATCC)	
Address of depository institution (including postal code and country)  10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit     December 30, 1999	Accession Number     PTA-1141
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-449  In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

***Escherichia coli* Plasmid TG-449****AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**CANADA**

The applicant hereby requests that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the furnishing of a sample of deposited biological material referred to in the application only be effected to an independent expert nominated by the Commissioner of Patents.

**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent office or any person approved by the applicant in the individual case.

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Registration), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the National Board of Patents and Registration or any person approved by the applicant in the individual case.

**ICELAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Icelandic Patent Office), or has been finally decided upon by the Icelandic Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

***Escherichia coli Plasmid TG-449*****NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in Rule 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**NORWAY**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Norwegian Patent office or any person approved by the applicant in the individual case.

**SINGAPORE**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent office or any person approved by the applicant in the individual case.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution American Type Culture Collection (ATCC)	
Address of depository institution <i>(including postal code and country)</i> 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit    December 30, 1999	Accession Number    PTA-1137
<b>C. ADDITIONAL INDICATIONS</b> <i>(leave blank if not applicable)</i> <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-376	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> <i>(if the indications are not for all designated States)</i>	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the international Bureau later <i>(specify the general nature of the indications, e.g., "Accession Number of Deposit")</i>	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on <u>page 12, line 20</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depository institution American Type Culture Collection (ATCC)	
Address of depository institution (including postal code and country)  10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit     December 30, 1999	Accession Number     PTA-1137
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<i>Escherichia coli</i> plasmid TG-376  In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

***Escherichia coli* Plasmid TG-376****AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**CANADA**

The applicant hereby requests that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the furnishing of a sample of deposited biological material referred to in the application only be effected to an independent expert nominated by the Commissioner of Patents.

**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent office or any person approved by the applicant in the individual case.

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Registration), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the National Board of Patents and Registration or any person approved by the applicant in the individual case.

**ICELAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Icelandic Patent Office), or has been finally decided upon by the Icelandic Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

***Escherichia coli Plasmid TG-376*****NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in Rule 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**NORWAY**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Norwegian Patent office or any person approved by the applicant in the individual case.

**SINGAPORE**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent office or any person approved by the applicant in the individual case.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

2. Statement Concerning Non-Prejudicial Disclosures or Exceptions to Lack of Novelty

Due to the publication of an abstract prior to a scientific meeting in November 1998, the applicants respectfully request that the subject international application be granted the respective provisions under national laws concerning Exceptions to Lack of Novelty in each of the designated countries. This is not an admission that the subject invention is not novel. Exception to lack of novelty is hereby requested for purposes of disclosure and precautionary measures.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/27862

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : C12N 1/21, 5/10, 15/12, 15/63

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 252.3, 254.11, 320.1; 530/350, 324, 329, 330; 536/23.5, 24.31

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE

search terms: parathyroid hormone receptor, PTH

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,494,806 A (SEGRE et al.) 27 February 1996, Figures 3A-3E; Figure 21; SEQ ID NO:3; column 19, line 50, through column 21, line 27.	1, 4, 7-9, 11-17

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

10 MAY 1999

Date of mailing of the international search report

20 MAY 1999

Name and mailing address of the ISA:US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DAVID ROMEIO

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/27862

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons.

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-17

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/27862

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/69.1, 252.3, 254.11, 320.1; 530/350, 324, 329, 330; 536/23.5, 24.31

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

Group I, claim(s) 1-17, drawn to a polynucleotide encoding a rδNt receptor, a rδNt receptor polypeptide and corresponding vectors and host cells.

Group II, claim(s) 18, drawn to an antibody that binds the rδNt receptor.

Group III, claim(s) 19, drawn to a method of screening for agonists and antagonists of PTH receptor activity.

The inventions listed as Groups I-III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The special technical feature of Group I is polynucleotide encoding a truncated form of a PTH receptor. Group II does not share this special technical feature because it is drawn to a product which is materially unlike the product(s) of Group I. Since the rδMt receptor is a truncated PTH receptor, the antibodies of Group II could be made alternatively with the full length receptor. The method of Group III does not require the products of either Group I or Group II and could alternatively practiced with a full length PTH receptor.

Groups II and III do not share a special technical feature because the product of Group II is not required to practice the method of Group III.